

**First Professional BUMS Degree Supplementary Examinations
December 2024
Tashreeh UI-Badan - Paper II
(2022 Scheme)**

Time: 3 hrs

Max Marks: 100

- Answer all questions to the point neatly and legibly • Do not leave any blank pages between answers • Indicate the question number correctly for the answer in the margin space
- Answer all parts of a single question together • Leave sufficient space between answers • Draw table/diagrams/flow charts wherever necessary

1. Multiple Choice Questions (20x1=20)

The Answers to MCQ questions (Q.No. i to Q.No. xx) shall be written continuously on the first two writing sheets (ie Page No. 3 & 4) only

- i. Which of the following is TRUE for Abdominal Aorta
 - a) It terminates in front of L4 Vertebra into right and left Common Iliac artery.
 - b) Sacral artery is the lateral branch.
 - c) Passes into Abdomen behind the diaphragm at the level of T10 vertebra.
 - d) Has renal arteries arising at the level of the body of L1 vertebra.
- ii. The Subpubic angle in female is
 - a) 60 degree
 - b) 90 degree
 - c) 110 degree
 - d) 15 degree
- iii. Sacral Hiatus transmits the following EXCEPT
 - a) Filum terminale
 - b) 5th pair of sacral nerve
 - c) One pair of Coccygeal nerve
 - d) 4th pair of sacral nerve
- iv. Duct of Bellini is present in
 - a) Malpighian corpuscles
 - b) Collecting tubules
 - c) Loop of Henle
 - d) Proximal convoluted tubule
- v. Tube carrying Spermatozoa from epididymis to the ejaculatory duct
 - a) Seminal vesicles
 - b) Urethra
 - c) Vas deferens
 - d) Vasa efferentia
- vi. The angle of Femoral torsion is
 - a) 15 degree
 - b) 125 degree
 - c) 90 degree
 - d) 50 degree
- vii. The greater sac and the Lesser sac communicates through
 - a) Duct of Bellini
 - b) Epiploic foramen
 - c) Obturator foramen
 - d) Foramen Caecum
- viii. Which of the following is an example of Uniaxial joint
 - a) Ankle joint
 - b) Knee joint
 - c) Wrist joint
 - d) Hip joint
- ix. Compression of Causes Tarsal Tunnel Syndrome
 - a) Peroneal nerve
 - b) Tibial nerve
 - c) Femoral nerve
 - d) Saphenous nerve
- x. Left gastric artery is a direct branch of
 - a) Common hepatic artery
 - b) Proper hepatic artery
 - c) Splenic artery
 - d) Coeliac trunk
- xi. "Most abundant Peyer's patches are present in"
 - a) Duodenum
 - b) Jejunum
 - c) Ileum
 - d) Caecum

(PTO)

- Short Answer Questions** **(8x5=40)**

- Long Answer Questions** (4x10=40)

- *****